

Degree Sum Conditions and Vertex-Disjoint Cycles in a Graph

Shinya Fujita

Hajime Matsumura

Department of Mathematics

Keio University

Yokohama 223-8522, Japan

shinyaa@comb.math.keio.ac.jp

musimaru@comb.math.keio.ac.jp

Masao Tsugaki

Department of Mathematical Information Science

Tokyo University of Science

Tokyo 162-8601, Japan

tsugaki@hotmail.com

Tomoki Yamashita

Department of Mathematics

School of Dentistry, Asahi University

Gifu 501-0296, Japan

tomoki@dent.asahi-u.ac.jp

Abstract

We consider degree sum conditions and the existence of vertex-disjoint cycles in a graph. In this paper, we prove the following: Suppose that G is a graph of order at least $3k + 2$ and $\sigma_3(G) \geq 6k - 2$, where $k \geq 2$. Then G contains k vertex-disjoint cycles. The degree and order conditions are sharp.

Keywords: vertex-disjoint cycle, degree sum condition

1 Introduction

We will generally follow notation and terminology of [1]. Let G be a simple graph. For a vertex x of a graph G , the neighborhood of x in G is denoted by $N_G(x)$, and $d_G(x) = |N_G(x)|$ is the degree of x in G . For a subgraph H of G and a vertex $x \in V(G) - V(H)$, we also denote $N_H(x) = N_G(x) \cap V(H)$ and $d_H(x) = |N_H(x)|$. For a subgraph H and a subset S of $V(G)$, $d_H(S) = \sum_{x \in S} d_H(x)$, the subgraph induced by S is denoted by $\langle S \rangle$, and $G - S = \langle V(G) - S \rangle$. We often identify an induced subgraph with its vertex set. For a graph G , $|G| = |V(G)|$ is the order of G , $\omega(G)$ is the number of components of G , $\delta(G)$ is the minimum degree of G , $\alpha(G)$ is the independence number of G and

$$\sigma_k(G) = \min \left\{ \sum_{x \in S} d_G(x) : S \text{ is an independent set of } G \text{ with } |S| = k. \right\}$$

(When $\alpha(G) < k$, we define $\sigma_k(G) = \infty$.)

For $X, Y \subseteq V(G)$, $E(X, Y)$ denote the set of edges of G joining a vertex in X and a vertex in Y . If $X = \{x\}$, we denote $E(x, Y)$ instead of $E(\{x\}, Y)$.

K_n denotes a complete graph of order n and P_n denotes a path of order n . For graphs G and H , $G \cup H$ denotes the union of G and H , and $G + H$ denotes the join of G and H . For a graph G , mG denotes the union of m copies of G . If a graph G is isomorphic to a graph H , we denote $G \simeq H$.

A forest is a graph each of whose components is a tree. A leaf is a vertex of a forest whose degree is at most 1.

In this paper, we consider degree sum conditions and the existence of vertex-disjoint cycles. The classical result of this problem was proved by Corrádi and Hajnal.

Theorem 1 (Corrádi and Hajnal [2]) *Suppose that $|G| \geq 3k$ and $\delta(G) \geq 2k$. Then G contains k vertex-disjoint cycles.*

Justesen improved Theorem 1 as follows.

Theorem 2 (Justesen [4]) *Suppose that $|G| \geq 3k$ and $\sigma_2(G) \geq 4k$. Then G contains k vertex-disjoint cycles.*

The degree condition in Theorem 2 is not sharp. Later, Enomoto and Wang independently improved Theorem 2 and got a sharp degree bound.

Theorem 3 (Enomoto [3], Wang [5]) *Suppose that $|G| \geq 3k$ and $\sigma_2(G) \geq 4k - 1$. Then G contains k vertex-disjoint cycles.*

Since $G_0 = K_{2k-1} + mK_1$ does not contain k vertex-disjoint cycles, and $\delta(G_0) = 2k - 1$ and $\sigma_2(G_0) = 4k - 2$, the degree conditions in Theorems 1 and 3 are weakest possible.

In this paper, we prove the following theorem.

Theorem 4 *Suppose that $k \geq 2$, $|G| \geq 3k + 2$ and $\sigma_3(G) \geq 6k - 2$. Then G contains k vertex-disjoint cycles.*

The sharpness of the degree condition is also shown by the graph G_0 since $\sigma_3(G_0) = 6k - 3$.

$K_{3k-1} \cup K_i$ ($i = 1, 2$) satisfies the degree condition of Theorem 4 since the independence number of this graph is 2, but does not contain k vertex-disjoint cycles. Hence $|G| \geq 3k + 2$ is also weakest possible.

Suppose that $n \geq 5$. Then $\sigma_3(P_n) = 4 = 6 \times 1 - 2$ but P_n does not contain a cycle. Hence $k \geq 2$ is necessary.

Note that the degree condition of Theorem 4 is weaker than those of Theorems 1 and 3:

If $\delta(G) \geq 2k$, then it is easy to see that $\sigma_2(G) \geq 4k - 1$ and $\sigma_3(G) \geq 6k - 2$. Suppose that $\sigma_2(G) \geq 4k - 1$. If we take three independent vertices x_1, x_2 and x_3 of G , then $d_G(x_1) + d_G(x_2) \geq 4k - 1$, $d_G(x_2) + d_G(x_3) \geq 4k - 1$ and $d_G(x_3) + d_G(x_1) \geq 4k - 1$. Hence we have $2(d_G(x_1) + d_G(x_2) + d_G(x_3)) \geq 12k - 3$, and $d_G(x_1) + d_G(x_2) + d_G(x_3) \geq 6k - 3/2$. This implies that $\sigma_3(G) \geq 6k - 2$.

Before proving Theorem 4, we will give some definitions.

Suppose that C_1, \dots, C_r are r vertex-disjoint cycles of a graph G . If C'_1, \dots, C'_r are r vertex-disjoint cycles of G and $|\bigcup_{i=1}^r V(C'_i)| < |\bigcup_{i=1}^r V(C_i)|$, then we call C'_1, \dots, C'_r

are shorter cycles than C_1, \dots, C_r . We call $\{C_1, \dots, C_r\}$ is minimal if G does not contain r vertex-disjoint cycles C'_1, \dots, C'_r such that $|\bigcup_{i=1}^r V(C'_i)| < |\bigcup_{i=1}^r V(C_i)|$. We call a cycle of order 3 a *triangle*.

We will use $C[u, v]$ to denote the segment of the cycle C from u to v (including u and v) under some orientation of C , and $C[u, v) = C[u, v] - \{v\}$ and $C(u, v) = C[u, v] - \{u, v\}$. Given a cycle C with an orientation, we let v^+ (resp. v^-) denote the successor (resp. the predecessor) of v along C according to this orientation. Analogously, $v^{2+} = (v^+)^+, v^{3+}, v^{2-} = (v^-)^-, v^{3-}, \dots$ are defined.

2 Proof of Theorem 4

The following lemmas will be used several times in this section.

Lemma 1 *Let r be a positive integer and C_1, \dots, C_r be r minimal vertex-disjoint cycles of a graph G . Then $d_{C_i}(x) \leq 3$ for any $x \in V(G) - \bigcup_{j=1}^r V(C_j)$ and for any i , $1 \leq i \leq r$. Furthermore, $d_{C_i}(x) = 3$ implies $|C_i| = 3$ and $d_{C_i}(x) = 2$ implies $|C_i| \leq 4$.*

Proof. This is easily seen by the minimality of $\{C_1, \dots, C_r\}$ ■

Lemma 2 *Suppose that F is a forest with at least two components and C is a triangle. Let x_1, x_2 and x_3 be leaves of F from at least two components. If $d_C(\{x_1, x_2, x_3\}) \geq 7$, then there are two vertex-disjoint cycles in $\langle F \cup C \rangle$ or there exists a triangle C' in $\langle F \cup C \rangle$ such that $\omega(\langle F \cup C \rangle - C') < \omega(F)$.*

Proof. Let $C = v_1v_2v_3v_1$ and F_1, F_2 and F_3 be components of F . Suppose that $x_1, x_2 \in V(F_1)$ and $x_3 \in V(F_2)$. If $d_C(x_1) = 3$, then $d_C(\{x_2, x_3\}) \geq 4$ and $N_C(x_2) \cap N_C(x_3) \neq \emptyset$. Hence we may assume that $v_3 \in N_C(x_2) \cap N_C(x_3)$. Then $C' = x_1v_1v_2x_1$ is a triangle such that $\omega(\langle F \cup C \rangle - C') < \omega(F)$. If $d_C(x_3) = 3$, then $d_C(\{x_1, x_2\}) \geq 4$ and $N_C(x_1) \cap N_C(x_2) \neq \emptyset$. Hence we may assume that $v_3 \in N_C(x_1) \cap N_C(x_2)$. Then $x_3v_1v_2x_3$ and $v_3P_{F_1}[x_1, x_2]v_3$ are two vertex-disjoint cycles, where $P_{F_1}[x_1, x_2]$ is the unique path in F_1 connecting x_1 and x_2 .

Next, suppose that $x_1 \in V(F_1)$, $x_2 \in V(F_2)$ and $x_3 \in V(F_3)$. We may assume that $d_C(x_1) = 3$ and $v_3 \in N_C(x_2) \cap N_C(x_3)$. Then $C' = x_1v_1v_2x_1$ is a triangle such

that $\omega(\langle F \cup C \rangle - C') < \omega(F)$. ■

Lemma 3 *Let C be a cycle and X be a set of three independent vertices. Suppose that $\langle C \cup X \rangle$ does not contain a cycle C' such that $|C'| < |C|$. If $|E(C, X)| \geq 7$, then $|C| = 3$, and $\langle C \cup X \rangle$ can be partitioned into a vertex-disjoint triangle and a path of order 3 connecting two vertices of X .*

Proof. Since $|E(C, X)| \geq 7$, $d_C(x) \geq 3$ for some $x \in X$. This implies that $|C| = 3$ by Lemma 1. Let $C = v_1v_2v_3v_1$ and $X = \{x_1, x_2, x_3\}$. We may assume that $d_C(x_1) = 3$. Since $d_C(\{x_2, x_3\}) \geq 4$, $N_C(x_2) \cap N_C(x_3) \neq \emptyset$. Without loss of generality, we may assume that $v_1 \in N_C(x_2) \cap N_C(x_3)$. Then $\langle C \cup X \rangle$ is partitioned into a triangle $x_1v_2v_3x_1$ and a path of order 3 $x_2v_1x_3$. ■

Lemma 4 *Let C be a cycle and T be a tree with three leaves x_1, x_2 and x_3 . If $d_C(\{x_1, x_2, x_3\}) \geq 7$, then there exists a cycle C' in $\langle C \cup T \rangle$ such that $|V(C')| < |V(C)|$, or $\langle C \cup T \rangle$ contains two vertex-disjoint cycles.*

Proof. This is immediate by Lemma 3. ■

Lemma 5 *Let G be a graph satisfying the assumption of Theorem 4 and C_1, \dots, C_{k-1} be $k-1$ minimal vertex-disjoint cycles of G . Suppose that there exists a tree T with at least three leaves, which is a component of $G - \bigcup_{i=1}^{k-1} V(C_i)$. Then G contains k vertex-disjoint cycles.*

Proof. Let $L = \bigcup_{i=1}^{k-1} V(C_i)$ and $X = \{x_1, x_2, x_3\}$ be a set of leaves of T . Since X is independent and $d_T(x) = 1$ for all $x \in X$, $d_L(X) \geq 6k - 2 - 3 = 6k - 5 > 6(k-1)$. Hence $d_{C_i}(X) \geq 7$ for some i , $1 \leq i \leq k-1$. By Lemma 4, there exist two vertex-disjoint cycles in $\langle X \cup C_i \rangle$ since $\{C_1, \dots, C_{k-1}\}$ is minimal. Hence we have k vertex-disjoint cycles of G . ■

Lemma 6 *Let G be a graph satisfying the assumption of Theorem 4 and C_1, \dots, C_{k-1} be $k-1$ minimal vertex-disjoint cycles of G . Suppose that $|G - \bigcup_{i=1}^{k-1} V(C_i)| = 4$ and $G - \bigcup_{i=1}^{k-1} V(C_i)$ is not connected and is not isomorphic to $2K_2$. Then there exist $k-1$ minimal vertex-disjoint cycles C'_1, \dots, C'_{k-1} such that $G - \bigcup_{i=1}^{k-1} V(C'_i)$ is connected.*

Proof. Let $L = \bigcup_{i=1}^{k-1} V(C_i)$, $H = G - L$ and $V(H) = \{x_1, x_2, x_3, x_4\}$. We have to consider the following three cases;

- (i) $H \simeq P_3 \cup K_1$,
- (ii) $H \simeq K_2 \cup 2K_1$, and
- (iii) $H \simeq 4K_1$.

Without loss of generality, we may assume that $x_1x_2, x_2x_3 \in E(G)$ for (i), and $x_1x_2 \in E(G)$ for (ii). In each of three cases, $X = \{x_1, x_3, x_4\}$ is independent and $d_H(X) \leq 2$. Hence $d_L(X) \geq 6k - 2 - 2 = 6k - 4 > 6(k - 1)$ and this implies that $d_{C_i}(X) \geq 7$ for some i , $1 \leq i \leq k - 1$. Then by Lemma 3, we can take minimal vertex-disjoint cycles C'_1, \dots, C'_{k-1} such that $\omega(G - \bigcup_{i=1}^{k-1} V(C'_i)) < \omega(H)$. Moreover, $G - \bigcup_{i=1}^{k-1} V(C'_i)$ contains a path of order 3 connecting two vertices of X . Hence $G - \bigcup_{i=1}^{k-1} V(C'_i) \not\simeq 2K_2$. By repeating this argument, we can get a conclusion. ■

Proof of Theorem 4. Let G be an edge-maximal counterexample. Since a complete graph of order at least $3k+2$ contains k vertex-disjoint cycles, G is not complete. Let x and y be non-adjacent vertices of G . Then $G' = G + xy$, the graph obtained from G by adding the edge xy , is not a counterexample by the maximality of G . Hence G' contains k vertex-disjoint cycles C_1, \dots, C_k and without loss of generality, we may assume that $xy \in E(C_k)$. This means that G contains $k - 1$ vertex-disjoint cycles C_1, \dots, C_{k-1} such that $\sum_{i=1}^{k-1} |V(C_i)| \leq n - 3$. Let $L = \langle \bigcup_{i=1}^{k-1} V(C_i) \rangle$ and $H = G - L$. Take $k - 1$ minimal vertex-disjoint cycles C_1, \dots, C_{k-1} so that

$$\omega(H) \text{ is as small as possible.} \tag{1}$$

Claim 1 *Each component of H is a path.*

Proof. This is immediate by Lemma 5. ■

Claim 2 *H is connected, or $|H| = 4$ and $H \simeq 2K_2$*

Proof. Suppose that H is not connected.

If $|H| \geq 5$ and $\omega(H) \geq 3$, then we can take three leaves x_1, x_2 and x_3 from three different components. If $|H| \geq 5$ and $\omega(H) = 2$, then there exists a component H' of H such that $|H'| \geq 3$. Since H' is a path by Claim 1, we can take two leaves x_1, x_2

from H' , and take a leaf x_3 from another component. In each case, $X = \{x_1, x_2, x_3\}$ is independent and $d_H(X) \leq 3$. Hence $d_L(X) \geq 6k - 2 - 3 = 6k - 5 > 6(k - 1)$ and this means that $d_{C_i}(X) \geq 7$ for some i , $1 \leq i \leq k - 1$. Then $d_{C_i}(x) \geq 3$ for some $x \in X$ and $|C_i| = 3$ by Lemma 1. By Lemma 2, we have $k - 1$ minimal vertex-disjoint cycles C'_1, \dots, C'_{k-1} such that $\omega(G - \bigcup_{j=1}^{k-1} V(C'_j)) < \omega(H)$ because G does not contain k vertex-disjoint cycles. But this contradicts the choice of cycles (1).

If $|H| = 4$ and $H \not\cong 2K_2$, then we can get the conclusion by Lemma 6.

Hence we may assume that $|H| = 3$. Let x and y be non-adjacent vertices of G . Then $G + xy$ contains k vertex-disjoint cycles D_1, \dots, D_k . Without loss of generality, we may assume that $xy \in E(D_k)$. If $|D_k| \geq 4$, then $|\bigcup_{i=1}^{k-1} V(D_i)| < |L|$, but this contradicts the minimality of L . Hence $|D_k| = 3$. If $G - \bigcup_{i=1}^k V(D_i) \neq \emptyset$, then $|\bigcup_{i=1}^{k-1} V(D_i)| < |L|$ since $|G - \bigcup_{i=1}^{k-1} V(D_i)| \geq 4$. Therefore, $V(G) = \bigcup_{i=1}^k V(D_i)$, $\{D_1, \dots, D_{k-1}\}$ is minimal and $G - \bigcup_{i=1}^{k-1} V(D_i)$ is connected. By the choice of cycles (1), H is connected. ■

We distinguish two cases according to the value of $|H|$.

CASE 1 $|H| \geq 5$

By Claims 1 and 2, H is a path. Let $x_1x_2 \cdots x_l$, where $l = |H|$, and let $X = \{x_1, x_3, x_l\}$. Then X is independent.

Claim 3 $d_{C_i}(X) \leq 6$ for any i , $1 \leq i \leq k - 1$.

Proof. Suppose that $d_{C_i}(X) \geq 7$ for some i , $1 \leq i \leq k - 1$. Since $d_{C_i}(x) \geq 3$ for some $x \in X$, $|C_i| = 3$ by Lemma 1. Let $C_i = v_1v_2v_3v_1$.

Suppose that $d_{C_i}(x_1) = 3$. Since $d_{C_i}(\{x_3, x_l\}) \geq 4$, $N_{C_i}(x_3) \cap N_{C_i}(x_l) \neq \emptyset$ and we may assume that $v_3 \in N_{C_i}(x_3) \cap N_{C_i}(x_l)$. Then $x_1v_1v_2x_1$ and $v_3x_3x_4 \cdots x_lv_3$ are two vertex-disjoint cycles in $\langle H \cup C_i \rangle$, and we have k vertex-disjoint cycles of G , a contradiction.

Hence $d_{C_i}(x_1) \leq 2$. Similarly, we have $d_{C_i}(x_l) \leq 2$. This means that $d_{C_i}(x_3) = 3$ and $d_{C_i}(x_1) = d_{C_i}(x_l) = 2$.

Suppose that $N_{C_i}(x_1) \neq N_{C_i}(x_l)$. Without loss of generality, we may assume that $v_1 \in N_{C_i}(x_1)$ and $v_2, v_3 \in N_{C_i}(x_l)$. Then $x_1x_2x_3v_1x_1$ and $x_lv_2v_3x_l$ are two vertex-

disjoint cycles in $\langle H \cup C_i \rangle$, and we have k vertex-disjoint cycles of G , a contradiction. Hence we have $N_{C_i}(x_1) = N_{C_i}(x_l)$ and we may assume that $\{v_1, v_2\} = N_{C_i}(x_1)$. If we take $C'_i = x_1 v_1 v_2 x_1$ and $C'_j = C_j$ for $j \neq i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is a tree with three leaves x_2, x_l and v_3 since otherwise we can find two vertex-disjoint cycles in $\langle H \cup C_i \rangle$. By Lemma 5, we have k vertex-disjoint cycles of G , a contradiction. Hence the proof is completed. \blacksquare

By Claim 3, we have

$$d_L(X) \leq 6(k-1).$$

On the other hand, since $d_H(X) = 4$,

$$d_L(X) \geq 6k - 2 - 4 = 6(k-1).$$

Hence $d_L(X) = 6(k-1)$ and $d_{C_i}(X) = 6$ for all i , $1 \leq i \leq k-1$. By Lemma 1, we have $|C_i| \leq 4$ since $d_{C_i}(x) \geq 2$ for some $x \in X$.

Claim 4 $|C_i| = 3$ for all i , $1 \leq i \leq k-1$.

Proof. Suppose that $|C_i| = 4$ and let $C_i = v_1 v_2 v_3 v_4 v_1$. By Lemma 1, $d_{C_i}(x) = 2$ for all $x \in X$.

Suppose that $N_{C_i}(x_1) \neq N_{C_i}(x_3)$. Then we may assume that $N_{C_i}(x_1) = \{v_1, v_3\}$ and $N_{C_i}(x_3) = \{v_2, v_4\}$. Note that there do not exist two vertex-disjoint cycles in $\langle H \cup C_i \rangle$, since otherwise we have k vertex-disjoint cycles of G , a contradiction.

Take $C'_i = x_1 v_1 v_2 v_3 x_1$ and $C'_j = C_j$ for $j \neq i$. Then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is a tree with three leaves x_2, x_l and v_4 . By Lemma 5, we have k vertex-disjoint cycles of G , but this is a contradiction.

Hence $N_{C_i}(x_1) = N_{C_i}(x_3)$. Similarly, we have $N_{C_i}(x_3) = N_{C_i}(x_l)$. Without loss of generality, we may assume that $N_{C_i}(x) = \{v_1, v_3\}$ for all $x \in X$. Taking $C'_i = x_1 x_2 x_3 v_1 x_1$ and $C'_j = C_j$ for $j \neq i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is a tree with three leaves x_4, v_2 and v_4 . By Lemma 5, this is also a contradiction. \blacksquare

Claim 5 Let $x \in \{x_1, x_l\}$. If $A \subset N_{C_i}(x)$ and $|A| = 2$, then $N_{C_i}(x_3) \setminus A = \emptyset$.




Figure 1: Configurations Type 1 and Type 2

Proof. Let $C_i = v_1v_2v_3v_1$. Suppose that the claim does not hold, and let $x = x_1$. Without loss of generality, we may assume that $v_1, v_2 \in N_{C_i}(x_1)$ and $v_3 \in N_{C_i}(x_3)$.

Take $C'_i = x_1v_1v_2x_1$ and $C'_j = C_j$ for $j \neq i$. Then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is a tree with three leaves x_2, x_l and v_3 since there do not exist two vertex-disjoint cycles in $\langle H \cup C_i \rangle$. By Lemma 5, we have k vertex-disjoint cycles of G , a contradiction.

For the case $x = x_l$, we can prove similarly. ■

Claim 6 *There exist only two type of configurations between H and C_i for all i , $1 \leq i \leq k - 1$. (See Figure 1.)*

Proof. Suppose that $d_{C_i}(x_1) = 3$. By Claim 5, we have $d_{C_i}(x_3) = 0$ and $d_{C_i}(x_l) = 3$ since $d_{C_i}(X) = 6$. (This is Type 1.)

Next, suppose that $d_{C_i}(x_1) \leq 1$. Since $d_{C_i}(X) = 6$, $d_{C_i}(\{x_3, x_l\}) \geq 5$. But this contradicts Claim 5.

Finally, suppose that $d_{C_i}(x_1) = 2$. By Claim 5, we have $N_{C_i}(x_1) = N_{C_i}(x_3)$. Since $d_{C_i}(X) = 6$, we have $d_{C_i}(x_l) = 2$ and $N_{C_i}(x_l) = N_{C_i}(x_3)$. (This is Type 2.)

Hence the claim is proved. ■

In each configuration, we find that $d_{C_i}(x_2) = d_{C_i}(x_4) = 0$ for any i , $1 \leq i \leq k-1$, since otherwise we can find two vertex-disjoint cycles in $\langle H \cup C_i \rangle$. This means that $d_G(x_2) = d_G(x_4) = 2$.

Let $C_1 = v_1v_2v_3v_1$. Since $\{x_2, x_4, v_3\}$ is independent,

$$6k - 2 \leq d_G(\{x_2, x_4, v_3\}) \leq 2 + 2 + 3(k-2) + 4 = 3k + 2,$$

but this is a contradiction since $k \geq 2$. This completes the proof of CASE 1.

CASE 2 $|H| \leq 4$.

Let $V(H) = \{x_1, \dots, x_{|H|}\}$. By Claims 1 and 2, we may assume that $x_1x_2, x_2x_3 \in E(G)$ if $|H| = 3$ and that $x_1x_2, x_3x_4 \in E(G)$ if $|H| = 4$.

Claim 7 *There exists i , $1 \leq i \leq k-1$ such that $|C_i| \geq 4$ and $|E(y, C_j)| \leq 3$ for any $y \in V(C_i)$ and $j \neq i$.*

Proof. Since $|G| \geq 3k+2$ and $|H| \leq 4$, $|L| \geq 3k-2 > 3(k-1)$. Hence there exists i , $1 \leq i \leq k-1$ such that $|C_i| \geq 4$.

We define a directed graph $\vec{D} = (V(\vec{D}), E(\vec{D}))$ as follows:

$$\begin{aligned} V(\vec{D}) &= \{C_i : |C_i| \geq 4, 1 \leq i \leq k-1\} \\ E(\vec{D}) &= \{(C_i, C_j) : |E(y, C_j)| \geq 4 \text{ for some } y \in V(C_i) \text{ and } j \neq i\} \end{aligned}$$

Suppose that \vec{D} contains a directed cycle. Without loss of generality, we may assume that $(C_1, C_2), (C_2, C_3), \dots, (C_m, C_1) \in E(\vec{D})$, where $m \geq 2$. Take $y_i \in V(C_i)$ so that $|E(y_i, C_{i+1})| \geq 4$. (Hereafter in the proof of this claim, let $C_{m+1} = C_1$.) Then there exist $v_{i+1}, w_{i+1} \in N_{C_{i+1}}(y_i)$ such that $y_{i+1} \notin C_{i+1}[v_{i+1}, w_{i+1}]$ and $C_{i+1}(v_{i+1}, w_{i+1}) \cap N_{C_{i+1}}(y_i) = \emptyset$. For $1 \leq i \leq m$, we define new cycles as

$$C'_i = y_i C_{i+1}[v_{i+1}, w_{i+1}] y_i.$$

Then $|\bigcup_{i=1}^m V(C'_i)| < |\bigcup_{i=1}^m V(C_i)|$, but this contradicts the minimality of L since $V(C'_{i+1})$ misses at least one neighbor of $N_{C_{i+1}}(y_i)$ for each i , $1 \leq i \leq m$. Hence \vec{D}

does not contain a directed cycle and an endvertex of a directed path is a desired cycle. ■

Without loss of generality, we may assume that C_1 satisfies the property of Claim 7.

Claim 8 $|\{y \in V(C_1) : |E(y, C_j)| = 3\}| \leq 2$ for any j , $2 \leq j \leq k - 1$.

Proof. Suppose not. Without loss of generality, we may assume that $|E(y, C_j)| = 3$ for any $y \in \{y_1, y_2, y_3\} \subset V(C_1)$. Let $v_1, v_2, v_3 \in N_{C_j}(y_1)$ and suppose that v_1, v_2 and v_3 appear in this order in C_j .

Suppose that $y_2v_1 \notin E(G)$. If $|N_{C_j}(y_2) \cap C_j(v_1, v_3)| \geq 2$, we can find two shorter cycles than C_1 and C_j . Since $d_{C_j}(y_2) = 3$, we have $|N_{C_j}(y_2) \cap C_j[v_3, v_1]| \geq 2$. In this case, we also find two shorter cycles than C_1 and C_j since $v_2 \in N_{C_j}(y_1)$.

Hence $y_2v_1 \in E(G)$. By symmetry, $v_1, v_2, v_3 \in N_{C_j}(y)$ for $y \in \{y_2, y_3\}$. But we can find two shorter cycles than C_1 and C_j since $|C_1| \geq 4$. ■

Claim 9 $E(x_2, C_1) \neq \emptyset$.

Proof. Suppose that $E(x_2, C_1) = \emptyset$. Let $Y = \{y_1, y_2, y_3, y_4\} \subset V(C_1)$ and suppose that y_1, y_2, y_3 and y_4 appear in this order in C_1 .

Subclaim 9.1 $f = 2|E(x_2, C_i)| + |E(Y, C_i)| \leq 12$ for any i , $2 \leq i \leq k - 1$.

Proof. Suppose that $f \geq 13$ for some i , $2 \leq i \leq k - 1$. Since $|E(Y, C_i)| \leq 10$ by the choice of C_1 and Claim 8, $|E(x_2, C_i)| \geq 2$. On the other hand, $|E(x_2, C_i)| \leq 3$ by Lemma 1.

Case A $|E(x_2, C_i)| = 3$.

In this case, we have $|C_i| = 3$ by Lemma 1. Furthermore, we have $|E(v, Y)| \leq 2$ for any $v \in V(C_i)$, since otherwise we can find two shorter cycles than C_1 and C_i in $\langle H \cup C_1 \cup C_i \rangle$. Then

$$f \leq 2 \times 3 + 2 \times 3 = 12,$$

a contradiction.

Case B $|E(x_2, C_i)| = 2$.

In this case, $|C_i| \leq 4$ by Lemma 1. Since $f \geq 13$, we have

$$|E(Y, C_i)| \geq 9. \quad (2)$$

Hence $d_{C_i}(y) \geq 3$ for some $y \in Y$. Without loss of generality, we may assume that $d_{C_i}(y_1) \geq 3$. Moreover, $d_{C_i}(y_1) = 3$ by the choice of C_1 . Let $Y' = \{y_2, y_3, y_4\}$.

Case B.1 $|C_i| = 4$.

Let $C_i = v_1v_2v_3v_4v_1$. We may assume that $v_1, v_2, v_3 \in N_{C_i}(y_1)$. Then $|E(v_j, Y')| \leq 1$ for $j \in \{1, 3, 4\}$ and $|E(v_2, Y')| \leq 2$ since otherwise we can find two shorter cycles than C_1 and C_i in $\langle C_1 \cup C_i \rangle$. Hence

$$|E(Y, C_i)| = d_Y(\{v_1, v_2, v_3, v_4\}) \leq 2 + 3 + 2 + 1 = 8,$$

but this contradicts (2).

Case B.2 $|C_i| = 3$.

Let $C_i = v_1v_2v_3v_1$. In this case, $N_{C_i}(y_1) = \{v_1, v_2, v_3\}$ and we may assume that $N_{C_i}(x_2) = \{v_1, v_2\}$. Then $|E(v_j, Y')| \leq 2$ for $1 \leq j \leq 2$ and $|E(v_3, Y')| \leq 1$, since otherwise we can find two shorter cycles than C_1 and C_i in $\langle C_1 \cup C_i \cup \{x_2\} \rangle$. Hence

$$|E(Y, C_i)| = d_Y(\{v_1, v_2, v_3\}) \leq 3 + 3 + 2 = 8,$$

but this contradicts (2). Hence Subclaim 9.1 is proved. ■

Since each of $\{x_2, y_1, y_3\}$ and $\{x_2, y_2, y_4\}$ is independent,

$$\begin{aligned} 2(6k - 2) &\leq 2d_G(x_2) + d_G(Y) \\ &\leq 12(k - 2) + 4 + 8 + |E(Y, H - \{x_2\})| \end{aligned}$$

by Subclaim 9.1. Hence $|E(Y, H - \{x_2\})| \geq 8$. Since $|H| \leq 4$, $|E(Y, x)| \geq 3$ holds for some $x \in V(H) - \{x_2\}$, but this contradicts the minimality of L by Lemma 1. Hence the proof of Claim 9 is completed. ■

First, we consider the case $|H| = 4$ and $H \simeq 2K_2$.

CASE 2.1 $H \simeq 2K_2$.

By Claim 9, $E(x, C_1) \neq \emptyset$ for all $x \in V(H)$. Then it is easy to see that $|C_1| \leq 6$. Let $x_1y \in E(G)$ for $y \in V(C_1)$. We give an orientation to C_1 so that $x_2y^- \notin E(G)$ if it is possible. Since at least one of x_3 and x_4 is not adjacent to y^- , we may assume that $x_3y^- \notin E(G)$. Then $Z = \{x_1, x_3, y^-\}$ is independent. Let $H' = \langle H \cup C_1 \rangle$.

Claim 10 $|E(Z, C_i)| \leq 6$ for any i , $2 \leq i \leq k-1$.

Proof. Suppose that $|E(Z, C_i)| \geq 7$ for some i , $2 \leq i \leq k-1$. We consider the following two cases.

Case A $5 \leq |C_1| \leq 6$, or $|C_1| = 4$ and there exists a cycle of order 4 containing x_1x_2 in H' .

In this case, we may assume that $x_2y' \in E(G)$ for $y' = y^{(|C_1|-3)^+}$. If we take $C'_1 = x_1C_1[y, y']x_2x_1$ and $C'_j = C_j$ for $2 \leq j \leq k-1$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal. Note that y^- does not lie in C'_i . By Lemma 1, $d_{C'_j}(y^-) \leq 3$ for $2 \leq j \leq k-1$ and $d_{C'_j}(y^-) = 3$ implies $|C'_j| = 3$. Since $|E(Z, C_i)| \geq 7$, $d_{C_i}(z) \geq 3$ for some $z \in Z$, and we have $|C_i| = 3$. Let $C_i = v_1v_2v_3v_1$.

Suppose that $d_{C_i}(x_1) = 3$. Since $d_{C_i}(\{x_3, y^-\}) \geq 4$, $d_{C_i}(x_3) \geq 1$ and we may assume that $x_3v_3 \in E(G)$. Take $C'_i = x_1v_1v_2x_1$ and $C'_j = C_j$ for $j \neq i$. Then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is connected or $G - \bigcup_{j=1}^{k-1} V(C'_j) \simeq P_3 \cup K_1$. By Lemma 6, this contradicts the choice of cycles (1). Therefore, $d_{C_i}(x_1) \leq 2$. Similarly, we have $d_{C_i}(x_3) \leq 2$.

Hence $d_{C_i}(y^-) = 3$ and $d_{C_i}(x_1) = d_{C_i}(x_3) = 2$. Without loss of generality, we may assume that $x_3v_3 \in E(G)$. Taking $C'_1 = x_1C_1[y, y']x_2x_1$, $C'_i = y^-v_1v_2y^-$ and $C'_j = C_j$ for $j \neq 1, i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is connected, or $G - \bigcup_{j=1}^{k-1} V(C'_j) \simeq P_3 \cup K_1$. But this contradicts the choice of cycles (1) by Lemma 6.

Case B $|C_1| = 4$ and there exists no cycle of order 4 containing x_1x_2 in H' .

By symmetry, we may assume that there exists no cycle of order 4 containing x_3x_4 in H' . In this case, $x_2y^{2+} \in E(G)$ and $d_{C_1}(x) = 1$ for all $x \in V(H)$.

By the choice of C_1 , $d_{C_i}(y^-) \leq 3$ holds. Then $d_{C_i}(\{x_1, x_3\}) \geq 4$, and we have $d_{C_i}(x_1) \geq 2$ or $d_{C_i}(x_3) \geq 2$. By Lemma 1, we have $|C_i| \leq 4$.

Case B.1 $|C_i| = 4$.

Let $C_i = v_1v_2v_3v_4v_1$. By Lemma 1, $d_{C_i}(x_1) \leq 2$ and $d_{C_i}(x_3) \leq 2$. Hence we have $d_{C_i}(y^-) = 3$ and $d_{C_i}(x_1) = d_{C_i}(x_3) = 2$.

Since $d_{C_i}(y^-) = 3$ and $d_{C_i}(x_3) = 2$ without loss of generality, we may assume that $v_1, v_2, v_3 \in N_{C_i}(y^-)$ and $v_3 \in N_{C'_i}(x_3)$. If we take $C'_1 = x_1C_1[y, y^{2+}]x_2x_1$, $C'_i = y^-v_1v_2y^-$ and $C'_j = C_j$ for $j \neq 1, i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is connected. This also contradicts the choice of cycles (1).

Case B.2 $|C_i| = 3$.

Let $C_i = v_1v_2v_3v_1$. Since $d_{C_i}(y^-) \leq 3$, we have $d_{C_i}(\{x_1, x_3\}) \geq 4$. Suppose that $d_{C_i}(x_1) = 3$. Then $d_{C_i}(x_3) \geq 1$ and without loss of generality, we may assume that $v_3 \in N_{C_i}(x_3)$. If we take $C'_i = x_1v_1v_2x_1$ and $C'_j = C_j$ for $j \neq i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is connected, or $G - \bigcup_{j=1}^{k-1} V(C'_j) \simeq P_3 \cup K_1$. By Lemma 6, this contradicts the choice of cycles (1). Hence $d_{C_i}(x_1) \leq 2$. Similarly, we have $d_{C_i}(x_3) \leq 2$. Then $d_{C_i}(y^-) = 3$ and $d_{C_i}(x_1) = d_{C_i}(x_3) = 2$, and we may assume that $v_1, v_2 \in N_{C_i}(x_1)$. If $v_3 \in N_{C_i}(x_3) \cap N_{C_i}(x_4)$, then $C'_i = x_1v_1v_2x_1$, $C'_k = x_3x_4v_3x_3$ and $C'_j = C_j$ for $j \neq i$ are k vertex-disjoint cycles in G . Hence $v_3 \notin N_{C_i}(x_1) \cap N_{C_i}(x_3)$. If $x_2v_3 \notin E(G)$, then $C'_i = x_1v_1v_2x_1$ and $C'_j = C_j$ for $j \neq i$ are $k-1$ minimal vertex-disjoint cycles and $G - \bigcup_{j=1}^{k-1} V(C'_j) \simeq K_2 \cup 2K_1$ or $P_3 \cup K_1$ since $v_3 \notin N_{C_i}(x_1) \cap N_{C_i}(x_3)$. By Lemma 6, this contradicts the choice of cycles (1). Therefore, $x_2v_3 \in E(G)$. Since $d_{C_i}(y^-) = 3$, $y^-v_3 \in E(G)$. Furthermore, since there is no cycle of order 4 containing x_3x_4 in H' and the minimality of L , $E(\{x_3, x_4\}, \{y, y^+\}) \neq \emptyset$. If we take $C'_1 = x_2v_3y^-y^{2+}x_2$, $C'_i = x_1v_1v_2x_1$ and $C'_j = C_j$ for $j \neq 1, i$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal and $G - \bigcup_{j=1}^{k-1} V(C'_j)$ is connected. But this contradicts the choice of cycles (1). This completes the proof of Claim 10. ■

Claim 11 $|E(Z, H')| \leq 9$.

Proof. Suppose that $|E(Z, H')| \geq 10$. Since $d_{H'}(y^-) \leq 4$, we have $d_{H'}(\{x_1, x_3\}) \geq 6$. On the other hand, $d_{H'}(\{x_1, x_3\}) \leq 6$ since $d_{C_1}(x) \leq 2$ for $x \in \{x_1, x_3\}$. Hence $d_{H'}(\{x_1, x_3\}) = 6$ and $d_{H'}(y^-) = 4$. Especially, $d_{C_1}(x_1) = d_{C_1}(x_3) = 2$. Then we have $x_1y^{2+}, x_3y, x_3y^{2+} \in E(G)$. Since $d_{H'}(y^-) = 4$, we have $x_2y^-, x_4y^- \in E(G)$. By the choice of an orientation of C_1 , $x_2y^+ \in E(G)$. But this gives two vertex-disjoint cycles in H' , $x_1x_2y^+yx_1$ and $x_3x_4y^-y^{2+}x_3$, a contradiction. \blacksquare

By Claims 10 and 11, we have

$$6k - 2 \leq d_G(Z) \leq 6(k - 2) + 9 = 6k - 3,$$

a contradiction. This completes the proof of CASE 2.1.

In the following, we consider the case $3 \leq |H| \leq 4$ and H is connected. Other than the assumptions we put at the beginning of Case 2, we may further assume that $x_2x_3 \in E(G)$ if $|H| = 4$. By Claim 9, we may assume that $x_2y \in E(G)$ for some $y \in V(C_1)$. Let $X = \{x_1, x_2, x_3\}$ and $Y = \{y^-, y, y^+\}$.

Claim 12 $g = |E(X \cup Y, C_i)| \leq 12$ for any i , $2 \leq i \leq k - 1$.

Proof. Suppose that $g \geq 13$ for some i , $2 \leq i \leq k - 1$. Then we have $|E(X, C_i)| \geq 5$ since $|E(Y, C_i)| \leq 8$ holds by the choice of C_1 and Claim 8. Hence $d_{C_i}(x) \geq 2$ for some $x \in X$ and we have $|C_i| \leq 4$ by Lemma 1.

Subclaim 12.1 $|E(Y, C_i)| \leq 7$ if $|C_i| = 4$.

Proof. Suppose that $|C_i| = 4$ and $|E(Y, C_i)| \geq 8$. Then $d_{C_i}(y') \geq 3$ for some $y' \in Y$. On the other hand, $d_{C_i}(y') \leq 3$ for any $y' \in Y$ by the choice of C_1 . Hence $d_{C_i}(y') = 3$ for some $y' \in Y$. Let $C_i = v_1v_2v_3v_4v_1$.

Suppose that $d_{C_i}(y^-) = 3$. Without loss of generality, we may assume that $v_1, v_2, v_3 \in N_{C_i}(y^-)$. Since $|E(Y, C_i)| \geq 8$, we have $|E(\{y, y^+\}, C_i)| \geq 5$ and $N_{C_i}(y) \cap N_{C_i}(y^+) \neq \emptyset$. But this implies that we can find two shorter cycles than C_1 and C_i in $\langle C_1 \cup C_i \rangle$, a contradiction.

Hence $d_{C_i}(y^-) \leq 2$. Similarly, we have $d_{C_i}(y^+) \leq 2$. But this means that $|E(Y, C_i)| \leq 7$, a contradiction. \blacksquare

Suppose that $|C_i| = 4$ and let $C_i = v_1v_2v_3v_4v_1$. Since $|E(Y, C_i)| \leq 7$ by Subclaim 12.1, we have $|E(X, C_i)| \geq 6$. On the other hand, $|E(X, C_i)| \leq 6$ holds by Lemma 1. Hence $|E(X, C_i)| = 6$ and $|E(Y, C_i)| = 7$. Without loss of generality, we may assume that $\{v_1, v_3\} = N_{C_i}(x_1) = N_{C_i}(x_3)$ and $\{v_2, v_4\} = N_{C_i}(x_2)$. Note that there exists a cycle of order 4 in $\langle (H \cup C_i) - \{v_j, v_{j+1}\} \rangle$ for any j , $1 \leq j \leq 3$. Since $|E(Y, C_i)| = 7$, $d_{C_i}(y') = 3$ for some $y' \in Y$ and $\{v_j, v_{j+1}\} \subseteq N_{C_i}(y')$ for some j , $1 \leq j \leq 3$. This means that we can find a triangle and a cycle of order 4 in $\langle H \cup C_1 \cup C_i \rangle$. This contradicts the minimality of L .

Hence we may assume that $|C_i| = 3$. Let $C_i = v_1v_2v_3v_1$ and $H'' = \langle H \cup C_1 \cup C_i \rangle$.

Suppose that $d_{C_i}(y^-) = 3$. Then $N_{C_i}(x_1) \cap N_{C_i}(x_2) = \emptyset$ and $N_{C_i}(x_2) \cap N_{C_i}(x_3) = \emptyset$, since otherwise we can find two vertex-disjoint triangles in H'' . Hence $|E(X, C_i)| \leq 6$. Also $N_{C_i}(y) \cap N_{C_i}(y^+) = \emptyset$, since otherwise we can find two vertex-disjoint triangles in $\langle C_1 \cup C_i \rangle$. Then $|E(\{y, y^+\}, C_i)| \leq 3$ and we get $g \leq 12$, a contradiction.

Hence $d_{C_i}(y^-) \leq 2$ and we have $d_{C_i}(y^+) \leq 2$, similarly. Furthermore, since we do not use the existence of the path $C_1[y^+, y^-]$ in the above argument, we have also $d_{C_i}(x_1) \leq 2$ and $d_{C_i}(x_3) \leq 2$ by the same argument. Therefore, $|E(\{x_1, x_3, y^-, y^+\}, C_i)| \leq 8$ and this implies that $|E(\{x_2, y\}, C_i)| \geq 5$.

Suppose that $d_{C_i}(y) = 3$. Since $|E(Y, C_i)| \leq 7$, we have $|E(X, C_i)| \geq 6$. Also, since $d_{C_i}(x_1) \leq 2$ and $d_{C_i}(x_3) \leq 2$, we have $d_{C_i}(x_2) \geq 2$ and this implies that $N_{C_i}(x_1) \cap N_{C_i}(x_2) \neq \emptyset$. Then we can find two vertex-disjoint triangles in H'' , a contradiction. Hence $d_{C_i}(y) \leq 2$. Again, we do not use the existence of the path $C_1[y^+, y^-]$, then we have $d_{C_i}(x_2) \leq 2$ by the same argument. But this means that $g \leq 12$, a contradiction. This completes the proof of Claim 12. ■

Since each of $\{x_1, x_3, y\}$ and $\{x_2, y^+, y^-\}$ is independent,

$$\begin{aligned} 2(6k - 2) &\leq d_G(X \cup Y) \\ &\leq 12(k - 2) + 10 + (|H| - 3) + |E(X, C_1)| + |E(Y, H)|. \end{aligned}$$

Hence

$$13 \leq |H| + |E(X, C_1)| + |E(Y, H)|. \quad (3)$$

We consider the following two cases.

CASE 2.2 $H \simeq P_4$.

By (3), $9 \leq |E(X, C_1)| + |E(Y, H)|$ and at least one of $|E(X, C_1)| \geq 5$ and $|E(Y, H)| \geq 5$ hold.

Let $H' = \langle H \cup C_1 \rangle$. Note that there is no triangle in H' by the minimality of L .

Claim 13 $|C_1| = 4$.

Proof. Suppose that $|E(X, C_1)| \geq 5$. Then $d_{C_1}(x) \geq 2$ for some $x \in X$ and we have $|C_1| = 4$ by Lemma 1.

Next, suppose that $|E(Y, H)| \geq 5$. This inequality implies that $d_Y(x) \geq 2$ for some $x \in H$ and also means that $|C_1| = 4$ by Lemma 1. ■

Let $C_1 = yy^+y'y^-y$. By symmetry of x_2 and x_3 , we have $E(x_3, C_1) \neq \emptyset$ by Claim 9.

Suppose that $x_3y' \in E(G)$. Then $d_{C_1}(x_2) = d_{C_1}(x_3) = 1$ since otherwise we can find a triangle in H' . If $x_1y^-, x_4y^+ \in E(G)$, then $x_1x_2yy^-x_1$ and $x_3x_4y^+y'x_3$ are two vertex-disjoint cycles in H' , and we have k vertex-disjoint cycles of G , a contradiction. If $x_1y^+, x_4y^- \in E(G)$, then $x_1x_2yy^+x_1$ and $x_3x_4y^-y'x_3$ are two vertex-disjoint cycles in H' . Hence $|E(G) \cap \{x_1y^-, x_4y^+\}| \leq 1$ and $|E(G) \cap \{x_1y^+, x_4y^-\}| \leq 1$. But this implies that $|E(X, C_1)| + |E(Y, H)| \leq 8$, a contradiction.

Hence $N_{C_1}(x_3) \subset \{y^-, y^+\}$. By symmetry of y^+ and y^- , we may assume that $x_3y^+ \in E(G)$. By replacing C_1 with $x_2x_3y^+yx_2$, we may assume that $\{x_1, x_4, y^-, y'\}$ induces P_4 . Since $x_1x_4 \notin E(G)$, we have either $\{x_1y^-, y'x_4\} \subset E(G)$ or $\{x_1y', y^-x_4\} \subset E(G)$. However, in the former case, $\langle H \cup C_1 \rangle$ has two vertex-disjoint cycles $x_1x_2yy^-x_1$ and $x_3x_4y'y^+x_3$, a contradiction. Thus, the latter case occurs. We have already seen $y'x_3 \notin E(G)$. By symmetry, we also have $x_2y^- \notin E(G)$. Then since $\langle H \cup C_1 \rangle$ has no triangle, we deduce $E(H, C_1) = \{x_1y', x_2y, x_3y^+, x_4y\}$. However, this implies that $|E(X, C_1)| + |E(Y, H)| \leq 6$. This is a contradiction and completes the proof of CASE 2.2.

CASE 2.3 $|H| = 3$




Figure 2: The configuration between H and C_1

By (3), we have $10 \leq |E(X, C_1)| + |E(Y, H)|$. Since there is no triangle in $\langle H \cup C_1 \rangle$ because of the minimality of L , $d_H(y) = 1$ and $d_H(y_0) \leq 2$ for $y_0 \in \{y^-, y^+\}$. Hence $|E(Y, H)| \leq 5$, and this implies that $|E(X, C_1)| \geq 5$. Then $|C_1| = 4$ by Lemma 1 since $d_{C_1}(x) \geq 2$ for some $x \in X$.

On the other hand, we have $|E(X, C_1)| \leq 6$ by Lemma 1. This implies that $|E(X, C_1)| + |E(Y, H)| \leq 11$.

Since each of $\{x_1, x_3, y\}$ and $\{x_2, y^-, y^+\}$ is independent,

$$\begin{aligned}
2(6k - 2) &\leq d_G(X \cup Y) \\
&\leq \sum_{i=2}^{k-1} |E(X \cup Y, C_i)| + 10 + |E(X, C_1)| + |E(Y, H)| \\
&\leq 12(k - 2) + 10 + 11 \\
&\leq 12k - 3
\end{aligned}$$

by Claim 12. Therefore,

$$11 \leq |E(X \cup Y, C_i)| \leq 12 \quad (4)$$

holds for any i , $2 \leq i \leq k - 1$.

Let $C_1 = yy^+y'y^-y$. Then we may assume that $\{x_1y^-, x_1y^+, x_2y, x_3y^-, x_3y^+\} \subset E(G)$ since $|E(X, C_1)| \geq 5$. (See Figure 2.) Let $Z = \{x_1, x_3, y\}$ and $Z' = \{x_2, y^-, y^+\}$.

Claim 14 $|E(Z, C_i)| \leq 6$ for any i , $2 \leq i \leq k - 1$.

Proof. Suppose that $|E(Z, C_i)| \geq 7$ for some i . If we take $C'_1 = x_1x_2x_3y^-x_1$ and $C'_j = C_j$ for $2 \leq j \leq k-1$, then $\{C'_1, \dots, C'_{k-1}\}$ is minimal. By Lemma 1, $d_{C'_j}(y) \leq 3$ for $2 \leq j \leq k-1$ and $d_{C'_j}(y) = 3$ implies $|C'_j| = 3$. Since $|E(Z, C_i)| \geq 7$ and $d_{C_i}(z) \geq 3$ for some $z \in Z$. Then $|C_i| = 3$, and let $C_i = v_1v_2v_3v_1$.

Suppose that $d_{C_i}(x_1) = 3$. If $d_{C_i}(y) = 2$, then $d_{C_i}(z') = 0$ for any $z' \in Z'$ since otherwise we can find two vertex-disjoint triangles in $\langle H \cup C_1 \cup C_i \rangle$. Then $|E(Z \cup Z', C_i)| = |E(X \cup Y, C_i)| \leq 9$, but this contradicts (4). Hence $d_{C_i}(y) \leq 1$, and we have $d_{C_i}(x_3) = 3$. In this case, we have also $d_{C_i}(z') = 0$ for any $z' \in Z'$ and this means that $|E(X \cup Y, C_i)| \leq 7$, a contradiction.

Hence $d_{C_1}(x_1) \leq 2$. Similarly, we have $d_{C_i}(x_3) \leq 2$. This means that $d_{C_i}(y) = 3$ and $d_{C_i}(x_1) = d_{C_i}(x_3) = 2$. In this case, we have $d_{C_i}(z') = 0$ for all $z' \in Z'$ again, and this implies that $|E(X \cup Y, C_i)| \leq 7$, a contradiction. This completes the proof of Claim 14. ■

Since Z is independent, we have

$$6k - 2 \leq d_G(Z) \leq 6(k - 2) + 9 = 6k - 3$$

by Claim 14, but this is a contradiction. This completes the proofs of CASE 2.3 and Theorem 4.

Acknowledgement

The authors would like to thank Professors Hikoe Enomoto, Yoshimi Egawa and Katsuhiko Ota for their valuable suggestions. Also, they would like to thank anonymous referee for valuable comments and helpful suggestions. This work was partially supported by the JSPS Research Fellowships for Young Scientists (to H. M.) and the 21st Century COE Program; Integrative Mathematical Sciences: Progress in Mathematics Motivated by Social and Natural Sciences (to T. Y.).

References

- [1] J. A. Bondy and U.S.R. Murty, Graph Theory with Applications, London Macmillan Elsevier 1976.
- [2] K. Corrádi and A. Hajnal, On the maximal number of independent circuits in a graph, Acta Math. Acad. Sci. Hungar. 14 (1963), 423–439.
- [3] H. Enomoto, On the existence of disjoint cycles in a graph, Combinatorica 18 (1998), 487–492.
- [4] P. Justesen, On independent circuits in finite graphs and a conjecture of Erdős and Pósa, Annals of Discrete Math. 41 (1989), 299–306
- [5] H. Wang, On the maximum number of independent cycles in a graph, Discrete Math. 205 (1999), 183–190.