

### 3 区別の発生状況

表3.1 スズメバチ類7種の延駆除件数 1983 - 2007

	コガタ	キイロ	ヒメ	モン	オオ	ヤド	クロ	計	コガタの占有率
千種	1575	28	21	71	37	1	4	1737	90.7%
東	336	1	2		3			342	98.2%
北	359		4	2	1			366	98.1%
西	235	1	5	1				242	97.1%
中村	196	1	4	1	1		1	204	96.1%
中	207	1	2	4	3			217	95.4%
昭和	777	4	10	10	12			813	95.6%
瑞穂	652		9	1	7			669	97.5%
熱田	180		1					181	99.4%
中川	194	2	5					201	96.5%
港	165		5	2				172	95.9%
南	420		8		4			432	97.2%
守山	1688	120	62	62	24			1956	86.3%
緑	1850	9	92	24	17		1	1993	92.8%
名東	1678	24	69	52	20	3	1	1847	90.9%
天白	1394	15	48	56	13			1526	91.3%
計	11906	206	347	286	142	4	7	12898	92.3%
種構成比	92.3%	1.6%	2.7%	2.2%	1.1%	0.0%	0.1%	100%	100%

コガタの占有率 総駆除件数にコガタスズメバチが占める割合  
 名東区では2000年に56個，2001年に74個の巣が保健所職員により駆除されている。

表3.2 コガタスズメバチの  
 最多駆除件数と営巣密度

区名	面積	最多 駆除件数	年度	営巣密度
千種	18.24	131	(2005)	7.2
東	7.72	35	(2004)	4.5
北	17.56	41	(2004)	2.3
西	17.90	27	(2004)	1.5
中村	16.32	24	(1997)	1.5
中	9.36	24	(2004)	2.6
昭和	10.93	83	(2005)	7.6
瑞穂	11.23	56	(2001)	5.0
熱田	8.16	23	(2001)	2.8
中川	32.01	29	(2001)	0.9
港	45.67	23	(1997)	0.5
南	18.47	44	(1997)	2.4
守山	33.99	177	(2004)	5.2
緑	37.85	170	(2005)	4.5
名東	19.42	155	(1998)	8.0
天白	21.62	152	(2004)	7.0
計	326.45	1059	(2004)	3.2

営巣密度 面積 1K<sup>2</sup>当たりの駆除件数

表3.3 スズメバチ類の区別駆除件数 千種区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983	10	2						12
1984	2							2
1985	12	1						13
1986	26							26
1987	16							16
1988	34	1						35
1989	21	2						23
1990	45	1	1					47
1991	57	3	2					62
1992	31	3	2	1		1		38
1993	25	5	2		1			33
1994	61	2		2	2			67
1995	116	3		3	1			123
1996	93			1	1			95
1997	85	1	4	4	4		2	100
1998	108		2	5				115
1999	102		3	4	4			113
2000	94		1	4	1			100
2001	125	1		6	1			133
2002	95			2	1			98
2003	104	1	2	7	3			117
2004	126		1	6	3			136
2005	131	2	1	16	10		1	161
2006	30			5	3		1	39
2007	26			5	2			33
計	1575	28	21	71	37	1	4	1737
種構成比	90.7%	1.6%	1.2%	4.1%	2.1%	0.1%	0.2%	100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.4 スズメバチ類の区別駆除件数 東区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983	3							3
1984	2							2
1985	4							4
1986	4							4
1987	8							8
1988	5							5
1989	2							2
1990								
1991	4							4
1992	2							2
1993	9							9
1994	11							11
1995	14							14
1996	17							17
1997	33	1						34
1998	29				1			30
1999	10							10
2000	18							18
2001	29		2		1			32
2002	26							26
2003	34				1			35
2004	35							35
2005	29							29
2006	2							2
2007	6							6
計	336	1	2		3			342
種構成比	98.2%	0.3%	0.6%		0.9%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.5 スズメバチ類の区別駆除件数

北区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤド	クロ	計
1983	2							2
1984								
1985	3							3
1986	1							1
1987	6		1					7
1988	9							9
1989	1							1
1990	6							6
1991	12							12
1992	4							4
1993	2							2
1994	6							6
1995	12							12
1996	27							27
1997	37		2					39
1998	32							32
1999	11							11
2000	12			1				13
2001	40		1					41
2002	29							29
2003	24							24
2004	41							41
2005	19			1				20
2006	13				1			14
2007	10							10
計	359		4	2	1			366
種構成比	98.1%		1.1%	0.5%	0.3%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.6 スズメバチ類の区別駆除件数

西区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤド	クロ	計
1983	1							1
1984								
1985	2	1						3
1986								
1987	1							1
1988	7							7
1989	3							3
1990	1							1
1991	11		1					12
1992	6							6
1993	2							2
1994	9			1				10
1995	17							17
1996	12							12
1997	25							25
1998	13		1					14
1999	4							4
2000	14							14
2001	26		1					27
2002	20							20
2003	19							19
2004	27							27
2005	9		1					10
2006	3							3
2007	3		1					4
計	235	1	5	1				242
種構成比	97.1%	0.4%	2.1%	0.4%				100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.7 スズメバチ類の区別駆除件数

中村区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヲバ	クロ	計
1983	2							2
1984	1							1
1985	1							1
1986								
1987	2							2
1988	2							2
1989								
1990	2							2
1991	3							3
1992	3							3
1993	5							5
1994	9	1						10
1995	7							7
1996	13							13
1997	24							24
1998	16				1			17
1999	5		2					7
2000	11						1	12
2001	18		1					19
2002	18			1				19
2003	14							14
2004	23							23
2005	7		1					8
2006	6							6
2007	4							4
計	196	1	4	1	1		1	204
種構成比	96.1%	0.5%	2.0%	0.5%	0.5%		0.5%	100%

表3.8 スズメバチ類の区別駆除件数

中区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヲバ	クロ	計
1983		1						1
1984	4							4
1985	1							1
1986	4							4
1987								
1988	4							4
1989	1							1
1990	6							6
1991	4							4
1992	3							3
1993	5							5
1994	14							14
1995	8			1				9
1996	12							12
1997	12							12
1998	16		1	2	1			20
1999	4		1	1				6
2000	8							8
2001	17				1			18
2002	18							18
2003	9							9
2004	22				1			23
2005	20							20
2006	8							8
2007	7							7
計	207	1	2	4	3			217
種構成比	95.4%	0.5%	0.9%	1.8%	1.4%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.9 スズメバチ類の区別駆除件数

昭和区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983	3							3
1984	8							8
1985	6							6
1986	10	2						12
1987	12		1	1				14
1988	16							16
1989	8							8
1990	19		2					21
1991	25							25
1992	7				1			8
1993	16		1	1				18
1994	37		1		1			39
1995	48			1				49
1996	37			1				38
1997	41			1				42
1998	53			1				54
1999	39		1		1			41
2000	60				1			61
2001	60		1					61
2002	46				2			48
2003	46		1	1	3			51
2004	77		1	1	1			80
2005	83		1	2	2			88
2006	11							11
2007	9	2						11
計	777	4	10	10	12			813
種構成比	95.6%	0.5%	1.2%	1.2%	1.5%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.10 スズメバチ類の区別駆除件数

瑞穂区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983	4							4
1984	12							12
1985	12		1					13
1986	15							15
1987	10		1					11
1988	15							15
1989	6							6
1990	18							18
1991	32		1					33
1992	7		1					8
1993	13							13
1994	42		1					43
1995	50				1			51
1996	36				1			37
1997	47				1			48
1998	54							54
1999	32		1					33
2000	25		2					27
2001	56							56
2002	40							40
2003	47				2			49
2004	25				1			26
2005	46		1	1	1			49
2006	2							2
2007	6							6
計	652		9	1	7			669
種構成比	97.5%		1.3%	0.1%	1.0%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.11 スズメバチ類の区別駆除件数

熱田区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヲバ	クロ	計
1983	1							1
1984	1							1
1985								
1986								
1987	2							2
1988								
1989	1							1
1990	1							1
1991	2							2
1992	3							3
1993	3							3
1994	6							6
1995	5							5
1996	2							2
1997	12							12
1998	17							17
1999	7							7
2000	13							13
2001	23		1					24
2002	20							20
2003	10							10
2004	15							15
2005	18							18
2006	9							9
2007	9							9
計	180		1					181
種構成比	99.4%		0.6%					100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.12 スズメバチ類の区別駆除件数

中川区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヲバ	クロ	計
1983								
1984	1							1
1985	1							1
1986								
1987	6	1						7
1988	6							6
1989	1							1
1990	1							1
1991	2							2
1992	8							8
1993	2							2
1994	4							4
1995	8							8
1996	15							15
1997	27							27
1998	13	1						14
1999	7		1					8
2000	15							15
2001	30							30
2002	8		1					9
2003	4		2					6
2004	9							9
2005	18		1					19
2006	3							3
2007	6							6
計	195	2	5					202
種構成比	96.5%	1.0%	2.5%					100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.13 スズメバチ類の区別駆除件数

港区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983	1							1
1984								
1985								
1986								
1987	3							3
1988	2		1					3
1989	1							1
1990								
1991	1							1
1992	6							6
1993	2							2
1994	3							3
1995	8							8
1996	11							11
1997	23		1					24
1998	11							11
1999	2							2
2000	13							13
2001	15		1					16
2002	14		1					15
2003	15		1	1				17
2004	14			1				15
2005	11							11
2006	7							7
2007	2							2
計	165		5	2				172
種構成比	95.9%		2.9%	1.2%				100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.14 スズメバチ類の区別駆除件数

南区

年度	コガタ	キイロ	ヒメ	モン	オオ	チャロ	クロ	計
1983								
1984	3							3
1985	3							3
1986	4							4
1987	8							8
1988	13							13
1989	2							2
1990	5							5
1991	12							12
1992	8							8
1993	9							9
1994	11							11
1995	20							20
1996	32							32
1997	44							44
1998	33		2					35
1999	9		1					10
2000	18		2		2			22
2001	27							27
2002	29							29
2003	33		2					35
2004	41							41
2005	40				1			41
2006	8				1			9
2007	8		1					9
計	420		8		4			432
種構成比	97.2%		1.9%		0.9%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.15 スズメバチ類の区別駆除件数

守山区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤロ	クロ	計
1983	1							1
1984	10	2						12
1985	3							3
1986	20	10		1				31
1987	23	1		1				25
1988	26	3	2					31
1989	9	3	1					13
1990	31	4	1	3				39
1991	66	8	1	3				78
1992	26	7	1	1	1			36
1993	27	4		1				32
1994	62	11		1	2			76
1995	94	4	2	3	1			104
1996	108	5		1				114
1997	92	1	5	4	1			103
1998	140	14	9	4	3			170
1999	67	9	13	3				92
2000	125	7	4	2	1			139
2001	137	4	3	5	3			152
2002	122		2	2	2			128
2003	130	8	3	8	4			153
2004	177	3	5	2	1			188
2005	146	12	9	15	4			186
2006	24		1		1			26
2007	22			2				24
計	1688	120	62	62	24			1956
種構成比	86.3%	6.1%	3.2%	3.2%	1.2%			100%

表3.16 スズメバチ類の区別駆除件数

緑区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤロ	クロ	計
1983	3							3
1984	10		1					11
1985	7							7
1986	7				1			8
1987	20		1					21
1988	26		2					28
1989	13							13
1990	42			1				43
1991	46		8					54
1992	32	2	1					35
1993	35	1	3				1	40
1994	123	3	3		1			130
1995	119		1	1				121
1996	105		1					106
1997	130	1	10	4				145
1998	128	1	15	4				148
1999	79		11	4	2			96
2000	98	1	5	1	1			106
2001	156				3			159
2002	132		8	1	2			143
2003	144		8	1				153
2004	167		6	1	1			175
2005	170		7	6	5			188
2006	26							26
2007	32		1		1			34
計	1850	9	92	24	17		1	1993
種構成比	92.8%	0.5%	4.6%	1.2%	0.9%		0.1%	100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上



表3.17 スズメバチ類の区別駆除件数

名東区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤド	クロ	計
1983	5							5
1984	1							1
1985	16	1	1					18
1986	33	1	2	1				37
1987	39	1		2				42
1988	64		1	2				67
1989	10							10
1990	37	2	1	2				42
1991	65	3	8	1	1			78
1992	41	1	5	1			1	49
1993	46	2	4	1	1			54
1994	90	3	2	3				98
1995	136	1	1	4	1			143
1996	104			3	1			108
1997	107	5	6	3				121
1998	155	1	3	10				169
1999	75	1	14	2	1			93
2000	72		3	2	2	1		80
2001	83		3	1	1	1		89
2002	116		4	1				121
2003	104		4	4	1			113
2004	108	1	2	3	2			116
2005	133	1	5	6	4	1		150
2006	23				3			26
2007	15				2			17
計	1678	24	69	52	20	3	1	1847
種構成比	90.9%	1.3%	3.7%	2.8%	1.1%	0.2%	0.1%	100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

表3.18 スズメバチ類の区別駆除件数

天白区

年度	コガタ	キイロ	ヒメ	モン	オオ	ヤド	クロ	計
1983	4							4
1984	3							3
1985	8				1			9
1986	24		1	1				26
1987	17							17
1988	36	1	4	2				43
1989	11							11
1990	46	2	4		1			53
1991	48	1	3					52
1992	29	3	1					33
1993	24	4	1	2				31
1994	68	2		2				72
1995	93		1	4				98
1996	71		3					74
1997	84		4	3				91
1998	77	1	3	7				88
1999	58		4	6	1			69
2000	113	1	4	5	2			125
2001	93		3	4	2			102
2002	85		2		1			88
2003	106		4	9	1			120
2004	152		2	4	1			159
2005	106		3	7	2			118
2006	19		1		1			21
2007	19							19
計	1394	15	48	56	13			1526
種構成比	91.3%	1.0%	3.1%	3.7%	0.9%			100%

\* 2006年・2007年は生活衛生センターによる駆除分のみを計上

図3.1 スズメバチ類の駆除件数の区別構成比 2002-2007




図3.2 コガタスズメバチの区別構成比 1983-2007




図3.3 キイロスズメバチの区別構成比 1983-2007




図3.4 ヒメスズメバチの区別構成比 1983-2007




図3.5 モンスズメバチの区別構成比 1983-2007




図3.6 オオスズメバチの区別構成比 1983-2007

